



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

STATISTICAL REPORTS OF MORBIDITY AND MORTALITY, STATES AND CITIES OF THE UNITED STATES (untabulated).

CONNECTICUT.—Month of May, 1911. Population of reporting towns, 1,110,617. Total number of deaths from all causes 1,421, including diphtheria 12, measles 12, scarlet fever 8, tuberculosis, pulmonary, 125, typhoid fever 4. Cases reported: Diphtheria, 148 in 37 towns, measles, 888 in 74 towns, scarlet fever, 320 in 60 towns, tuberculosis, pulmonary, 169 in 43 towns, typhoid fever, 27 in 17 towns.

FLORIDA.—Week ended June 3, 1911. Reports from the State board of health show diphtheria in 2 localities (Sanford and Tampa) with 3 cases, malaria in 7 localities with 33 cases, smallpox in 11 counties with 67 cases, tuberculosis in 7 localities with 11 cases, typhoid fever in 1 locality (Tampa) with 17 cases.

KENTUCKY—*Louisville*.—Month of April, 1911. Population, 223,928. Total number of deaths from all causes 311, including diphtheria 4, measles 4, tuberculosis, pulmonary, 52, typhoid fever 3. Cases reported: Diphtheria 14, measles 12, scarlet fever 34, tuberculosis, pulmonary, 50.

LOUISIANA—*Shreveport*.—Month of May, 1911. Population, 28,015. Total number of deaths from all causes 86, including tuberculosis 14, typhoid fever 2.

MICHIGAN.—Month of April, 1911. Population 2,810,173. Total number of deaths from all causes 3,362, including diphtheria 35, measles 31, scarlet fever 25, tuberculosis 264, typhoid fever 44.

NEW YORK.—Month of April, 1911. Population 9,113,614. Total number of deaths from all causes 13,330, including diphtheria 185, measles 143, scarlet fever 200, tuberculosis 1,549, typhoid fever 80. Cases reported: Diphtheria 1,613, measles 7,794, scarlet fever 3,700, smallpox 5, tuberculosis, pulmonary or laryngeal, 2,701, typhoid fever 339.

OKLAHOMA.—Month of April, 1911. Population 1,657,155. Total number of deaths from all causes 704, including diphtheria 3, scarlet fever 10, tuberculosis 52, typhoid fever 10. Cases reported: Diphtheria 36, scarlet fever 240, smallpox 219, tuberculosis 110, typhoid fever 73.

TEXAS—*Fort Worth*.—Month of May, 1911.—Population 73,312. Total number of deaths from all causes 106, including scarlet fever 1, tuberculosis 12, typhoid fever 6. Cases reported: Diphtheria 3, measles 4, scarlet fever 1, smallpox 1, tuberculosis 12, typhoid fever 12.